

Polarization enhancement in $\vec{d}(\vec{p}, ^2\text{He})n$ reaction: Nuclear teleportation

February 9, 2008

S. Hamieh

*Kernfysisch Versneller Instituut,
Zernikelaan 25, 9747 AA Groningen, The Netherlands.*

Abstract

I show that an experimental technique used in nuclear physics may be successfully applied to quantum teleportation (QT) of spin states of massive matter. A *new* non-local physical effect the ‘quantum-teleportation-effect’ is discovered for the nuclear polarization measurement. Enhancement of the neutron polarization is expected in the proposed experiment for QT that discriminates *only* one of the Bell states.

1 Introduction

Discovery of quantum teleportation (QT) [1] is one of the most profound results of quantum information theory. By means of a classical communication channel and a quantum source realized by a non-local entangled state such as an EPR-pair of particles, the teleportation process allows to transmit an unknown quantum state from a sender to a receiver which are spatially separated. Experimental realizations of QT have so far been limited to teleportation of light rather than massive fermion [2, 3]. The present work gives an experimental scheme for QT of proton spin to neutron spin in $\vec{d}(\vec{p}, ^2\text{He})n$ reaction.

2 QT with $\vec{d}(\vec{p}, ^2\text{He})n$ reaction

Preparing a polarized $\vec{d}$ target 3S_1 and $m_s = 0$ state with polarized proton beam and selecting knockout reaction, will lead to the QT of the beam polarization to the outgoing neutron. In fact, for clarity reason two assumptions are made:

Assumption I: assuming the initial states of the deuteron target and the proton beam are pure¹, under such condition, the deuteron spin state could be

¹General procedure of QT of mixed state through noisy channel could be found in Ref. [4].

written as

$$|\psi^+\rangle_{23} = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle), \quad (1)$$

and the proton spin state as

$$|\Phi\rangle_1 = a|0\rangle + b|1\rangle, \quad (2)$$

where 1, 2, 3 denote the two protons, and the neutron states, respectively, and $|0\rangle$, $|1\rangle$ correspond to the positive, negative projection of the spin to the quantization axis, respectively. Using the so-called Bell's basis such that

$$|\psi^\pm\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle); \quad |\phi^\pm\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle), \quad (3)$$

the initial spin state of the system $(\vec{d}, \vec{p})$ could be written as follows

$$\begin{aligned} |\Psi\rangle_{123} = & \frac{1}{2} \{ |\phi^+\rangle_{12}(a|1\rangle + b|0\rangle)_3 + |\phi^-\rangle_{12}(a|1\rangle - b|0\rangle)_3 + |\psi^+\rangle_{12}(a|0\rangle + b|1\rangle)_3 \\ & + |\psi^-\rangle_{12}(a|0\rangle - b|1\rangle)_3 \}. \end{aligned} \quad (4)$$

Assumption II: assuming that the polarization transfer and the induced polarization at 0° are neglected for knockout reaction. This assumption is supported by the low measured $K_y^{y'} \sim -0.1$ [5] of the reaction $1 + \frac{\vec{1}}{2} \rightarrow \frac{\vec{1}}{2} + 0$ that has the same spin structure of $\vec{d}(\vec{p}, ^2\text{He})n$ reaction for unpolarized d . This suggests that the $\vec{1} + \frac{\vec{1}}{2} \rightarrow \frac{\vec{1}}{2} + 0$ will have low $K_y^{y'}$. Therefore, measuring the polarization of the high energy neutron, i.e. low pp relative energy will force the two outgoing proton state to be singlet and project the wave function of Eq. 4 after measurement into [5]

$$|\Psi\rangle_{123} = |\psi^-\rangle_{12}(a|0\rangle - b|1\rangle)_3. \quad (5)$$

Clearly, the polarization of the proton target has been teleported to the outgoing neutron polarization. By a 180° rotation around the y direction the state of the proton beam will be recovered by the outgoing neutron. Note that this experiment is not complete teleportation because we discriminate only the $|\psi^-\rangle$ state².

If the teleportation is taking place in such reaction a strong correlation should be observed between the polarization of the outgoing neutron and the polarization of the proton beam. Specially, when the polarization of the beam is in x direction the expected polarization of the outgoing neutron is in $-x$ direction (flipped) however for y polarization no flipping should be observed. Special requirement of the experiment are listed in the table 1.

²Contamination from higher multipole is estimated to be 4% and it is neglected here.

Table 1: Special requirement for the QT experiment with $\vec{d}(\vec{p}, {}^2\text{He})n$ reaction

target polarization	beam polarization	E_d	reaction type	detection systems
$P_z \sim 0$	$a 0\rangle + b 1\rangle$	170 MeV	proton	P_{neutron}
$P_{zz} \sim -2$			knockout	E_{neutron}

3 Conclusion

I have shown that the teleportation of massive Fermion could be implemented using the available experimental technique. I have suggested a new physical effect for the nuclear polarization measurement namely the ‘quantum-teleportation-effect’. For the proposed experiment the expected neutron polarization is more or less zero, however taking into account such effect the neutron polarization should be enhanced and strongly correlated to the beam polarization. Dynamical study that couples the quantum entanglement with the nuclear decoherence due to the polarization transfer and the induced polarization will be addressed elsewhere.

References

- [1] Bennett C. H. *et al.*, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [2] Bauwmeester D. *et al.*, *Nature* **390**, 575 (1997).
- [3] Boschi D. *et al.*, *Phys. Rev. Lett.* **80**, 112, (1998).
- [4] S. Oh, S. Lee, H. Lee, *quant-ph/0206173*, accepted for publication in *Phys. Rev. A*.
- [5] Ohlsen G., *Rep. Prog. Phys.* **35**, 717 (1972).